

# Piggery Farming

1. Introduction to pig farming
2. Breeds of pigs
3. Breeds for North East India
4. Ghungroo Pig
5. Breeding Management
6. Care and management of piglets
7. Feeding Management of pigs
8. Housing management of pigs
9. Prevention and control of pig diseases
10. Related Resources

## Introduction to pig farming

### Advantages of pig rearing

- Pigs convert inedible feeds, forages, certain grain byproducts obtained from mills, meat by products, damaged feeds and garbage into valuable nutritious meat. Most of these feeds are either not edible or not very palatable to human beings
- Pig grows fast and is a prolific breeder, farrowing 10 to 12 piglets at a time. It is capable of producing two litters per year under optimal management conditions
- The carcass return is quite high i.e. 60-80 percent of live body weight
- With a small investment on building and equipment, proper feeding and sound disease control programme the farmer can profitably utilize his time and labour in this subsidiary occupation
- The faeces of pigs is used as a manure to maintain soil fertility

## Breeds of pigs

The indigenous pig has been the basis used for pig production for a long period of time. It is small in size. Improved breeds are now being used for grading up the form the basis for pig production in the rural areas.

The imported pig breeds reared in India are as follows

### Large white Yorkshire



- Most extensively used exotic breed in India
- Body colour is solid white with occasional black – pigmented spots
- Erect ears, snout of medium lengths and dished face

## For: M Sc IV Semester Zoology Paper 4 Economic Zoology

- Excellent breed for the purpose of cross breeding
- Prolific breeds
- Mature boar 300-400 k.g
- Mature sow 230-320 kg

### Landrace



- White with black skin spot
- Long body, large drooping ears and long snout
- Prolific breeding and efficient in utilizing feed
- Carcass quality equal to Yorkshire
- Excellent breed for crossbreeding
- Mature boars weigh 270-360 kg
- Mature sow weighs 200-320 kg

### Middle white Yorkshire



- Used in some areas in India
- Grows rapidly gives good dressing percentage
- Not so prolific as large white Yorkshire
- Males 250-340 kg
- Females 180-270 kg

### Breeds for North East India

Pig farming is very important component in North East India. Out of total pig population in India, 28% are grown in this region. Following are the commonly grown breeds. Good quality breeds are available at ICAR Research Complex for NEH Region, Barapani – 793 103, Meghalaya.

*By: Dr. Santosh Thakur, Lecturer, S.S. In Zoology and Biotechnology, Vikram University, Ujjain*

**For: M Sc IV Semester Zoology Paper 4 Economic Zoology**



**Hampshire**



**HSX 1**



**Large White Yorkshire**



**Duroc**

*By: Dr. Santosh Thakur, Lecturer, S.S. In Zoology and Biotechnology, Vikram University, Ujjain*



## Landrace



## Indigenous type

### Ghungroo Pig



Ghungroo an indigenous strain of pig first reported from North Bengal is popular among the local people because of high prolificacy and ability to sustain in low input system. This breed/strain produces high quality pork utilizing agricultural byproducts and kitchen wastes. Ghungroo are mostly black coloured with typical 'Bull dog' face appearance, with a litter size of 6-12 piglets, individually weigh about 1.0 kg at birth and 7.0 – 10.0 kg at weaning. Both sexes are very much docile and easy to handle. In the breeding tract they are maintained under scavenging system and mainly act as insurance to the rainfed agriculture.

## Breeding Management

Age to breed gilts	8 months
Weight of breed gilts	100-120 kg
Length of heat period	2-3 days

## For: M Sc IV Semester Zoology Paper 4 Economic Zoology

Best time to breed in heat period	Gilts – first day
	Sows- Second day
Number of services per sow	2 services at an interval of 12-14 hours
Period of oestrous cycle	18-24 days (Average 21 days)
Occurrence of heat after weaning	2-10 days
Gestation period	114 days

### Selection of breeding stock

Important characteristics which need to be considered in developing a good sow herd are

- Size of litters
- Strength and vigour of litters
- Milking ability
- Temperament

Gain and feed efficiency, fertility, selection of individual animals from a herd is more important than the selection of a particular breed. Each producer at the time of setting up his herd should purchase his animals from a reliable disease free herd and should obtain as much information of the animals as possible. Once the herd is established the selection of the gilts and boars for replacement in the breeding herd should be based on the types and performance.

### Selection of gilts (female)



- Selection of gilts for the breeding herd should be made at market weight ie. When the animals weigh about 90 kg
- Select gilts from sows which have consistently farrowed and weaned large litters
- They have reached market in minimum time and have desirable market type
- It would be desirable to choose gilts whose litter mates and other fullsibs have given good performance in daily weight gain and feed conversion efficiency

### Selection of boars (male)





## For: M Sc IV Semester Zoology Paper 4 Economic Zoology

- Selection of boar is extremely important, particularly for a small breeding farm or unit
- The boar should be purchased from a breeder or a farm maintaining adequate information on its performance
- The boar should be selected from a dam which has consistently farrowed and weaned high litters
- A good boar will weigh 90 kg in about 5-6 months will be of good type and will be strong on feet and legs
- The feed conversion from weaning to 90 kg weight would be the most desirable.

## Points to be considered while replacing boars and gilts

- The mother of the pig to be selected should have had large litters of 8 piglets or more. The weaning weight (at 56 days) of a litter in case of a gilt selection should have been 120 kg and in the case of sow it should not less than 150 kg
- The gilt or the boar should have reached a body weight of about 90 kg in about 6 months
- The pig should have adequate length and depth of the body, thick well muscled hams should be firm and trim
- The pig should have sound feet and legs
- The back fat prove in live animals is not yet being practiced in this country. For gilt back fat thickness of 4 cm or less, boars 3.2 or less
- Gilts should have a minimum of 12 evenly spaced, functional teats. An animal with blind teats should be avoided as there will be little or no milk from these teats and the defect is heritable
- Negative blood test for both brucellosis and leptospirosis should be made during selection and the pigs should be vaccinated against swine fever
- Pigs should be free from other diseases and physical defects

## Age of breeding stock

Well developed gilts may as a general rule bred to farrow when 12-14 months old. This depends more on development than on age. Gilts should weigh at least 100 kg before breeding. Ovulation rate increases during successive oestrous periods (up to fifth) following puberty. Thus it is advantageous to delay the breeding of gilts until the second or third oestrous. Litter size increases on an average in succeeding pregnancies up to 5<sup>th</sup> or 6<sup>th</sup> litter. It is therefore advantageous to cull the sow from a breeding herd or a commercial herd after her fifth or sixth litter as the litter size goes down thereafter.

## Detection of heat

The average length of oestrous cycle in pigs is 21 days. The oestrous symptoms last for five to seven days beginning with vulvar swelling and vaginal discharge. In true oestrous there is frequent urination, reduced appetite, mounting and standing for service detected by the erection of ears and immobility when normal pressure is applied to the back. The application of pressure on the back is used to determine the correct breeding time. Animals with a predisposition for weak oestrous should be brought near the boar to exhibit heat symptoms a little more clearly.

Best time for breeding is during the latter half of the first day or early on the second day of oestrous.

## Flushing

It is the method of feeding sows and gilts before breeding. A good grower ration fed to sows and gilts seven to ten days before breeding helps in increased ovulation rates in them. After breeding sows and gilts

## **For: M Sc IV Semester Zoology Paper 4 Economic Zoology**

should be fed a limited but well balanced ration until the last six weeks of pregnancy and then full feeding should be resumed.

### **Care and management of pregnant animals**

The gestation period of sow varies from 109-120 days with an average of 114 days. Pregnant animals should be housed in groups in separate enclosures and should not be mixed with new animals to avoid fighting which at times may result in abortion. It would also be advisable to house pregnant gilts and sows in separate groups during gestation. About 3 m<sup>2</sup> of dry housing should be available for each sow. The pregnant animals should be allowed to move about every day in the morning on a free range or a pasture if available. A pasture area is presumed to be clean if a cultivated crop was raised.

### **Management at farrowing**

Farrowing time is the critical time in pig production. Death rate is high during farrowing and the first week after farrowing. Sows may be farrowed in pens equipped with guard rails and a creep space in farrowing crates or in farrowing stalls. A pen equipped with guard rails and a creep space is adequate. The pen should be maintained at 24°C to 28°C until the piglets are three or four days old and at 18°C to 22°C until the piglets are approximately six weeks old. The heat lamps should be hung 45 cm from the floor and suitably protected. The farrowing pens should be thoroughly cleaned before the sow is brought in. This will prevent a large number of diseases of piglets. The sow should be watched closely for determining the approximate time of farrowing and feed should not be given 12 hours before farrowing.

### **Care during farrowing**

An attendant should be on hand when the sow farrows. Otherwise many piglets will die. It takes generally 2 to 4 hours for complete farrowing to take place. The piglets should be removed as they are farrowed and kept warm in the creep space until farrowing is complete. Each piglet should be cleaned of all mucus to ensure that the breathing passages are clear. The navel cord should be tied 2-5 cm away from the navel, cut with a disinfected pair of scissors and the stumps painted with iodine. Piglets should be allowed to be nursed after birth. In about 2 days they settle down to their individual teats. They nurse 8-10 times in 24 hours in the initial period. Trampling by the sow should be prevented during the first two weeks.

### **Care and management of piglets**

#### **Removal of needle teeth**

Piglets are born with four pairs of sharp teeth, with two pairs on each jaw. They are of no practical value to the piglets and they may irritate the sow's udder during nursing or cause injury to other piglets. Clipping of these teeth shortly after birth will prevent the injury of the udder caused by the needle teeth.

#### **Anaemia in piglets**

Anaemia is a common nutritional disease in piglets. This condition can be prevented and cured by supplying iron either orally or by injection. Oral administration consists of spraying or swabbing the sow's udder with a saturated solution of ferrous sulphate (0.5 kg of ferrous sulphate in 10 litres of hot water). This solution must be applied daily from birth until the piglets start eating creep feed. Intra muscular injection of iron –dextran compounds is the more effective method of preventing anaemia.

## **Raising orphan piglets**

The death of a sow after farrowing, mastitis, lactation failure of litters larger than the sow is able to raise, result in orphan pigs. If another sow has farrowed within a short time previously, the orphan piglets may be transferred to her. This transfer must be made within a few days after farrowing because those section of sow's udder are not used soon cease producing milk.

Orphan piglets can also be raised with milk replacer. Milk replacer consists of one egg yolk thoroughly mixed with one litre of cow milk.

## **Castration**

The male piglets not selected for breeding may be castrated when they are three to four weeks old.

## **Separation of piglets from mother (Weaning)**

Normal weaning age of piglets is at 8 weeks age.

## **Feeding Management of pigs**

### **Points to be considered while formulating feeding ration**

- Most economical ingredients should be selected
- Grains- maize, sorghum, oat, other millets, wheat and rice should form the basic ingredients
- Protein supplements - oil cakes and fishmeal and meat meal
- No vitamin supplements is necessary if the pigs are allowed to pasture or are fed fresh green legumes. Vitamin B 12 supplement would be necessary if little or no animal protein is fed
- Antibiotic supplements at the rate of 11 mg of antibiotic per kilogram of ration
- Mineral supplements should be provided

The following table gives specifically the various requirements in the formation of creep, grower and finisher rations for pigs.

Nutrients	Creep feed (Up to weaning)	Grower ration (20-40 kg)	Finisher ration (40-90 kg)
Protein supplement (%)			
Oilcakes	16-18	14-16	13-14
Animal protein	8-10	4	2
Grains (Maize, sorghum, millets or combination of grains) (%)	60-65	50-55	40-50
Wheat bran or rice bran (%)	5	10	20
Lucerne meal (%) if available	--	5-8	--
Mineral mixture(%)	0.5	0.5	0.5
Antibiotic supplement (mg)	40	20	10

### **The composition of the concentrate feed for various age groups pigs**

*By: Dr. Santosh Thakur, Lecturer, S.S. In Zoology and Biotechnology, Vikram University, Ujjain*



## For: M Sc IV Semester Zoology Paper 4 Economic Zoology

Ingredients	Creep (14 <sup>th</sup> to 56 <sup>th</sup> day feed)	Grower ration (up to 40 kg)	Finisher ration(40-90 k.g)	Pregnant and nursing sows
Maize or sorghum or broken wheat, broken rice and barley in convenient combinations	65	50	50	50
Oil cakes (groundnut oil cake, soya bean oil-cake, sesame oil cake, linseed oilcake)	14	18	20	20
Molasses	5	5	5	5
Wheat bran or rice bran	10	1.5	25	18
Fishmeal or meat meal or cooked offal, skim milk powder dairy wastes	5	5	3	5
Mineral mixture	1	1.5	1.5	1.5
Salt	--	0.5	0.5	0.5

The most convenient way to feed animals on a farm is to prepare the complete ration recommended for different classes and give the pigs the amount they will eat without waste two or three times daily. The following is the approximate amount of dry feed the pigs will consume.

Weight of pig (kg)	Daily consumption of feed (kg) per pig
25	2.0
50	3.2
100	5.3
150	6.8
200	7.5
250	8.3

All grains in mixed feeds should be ground. Generally feeding in the form of wet mash is not superior than (Slop feeding) dry feeding. Slop feeding requires more time and excessive labour. If a ration is fairly high in fibre, pelleting the feed may increase the rate and efficiency of gain in weight. Pelleting may also decrease the amount of feed that is wasted. It is important not to overfeed sows which have been bred. Over fat sows are apt to produce weak pigs and crush more piglets at farrowing. Sows should gain about 35 kg and gilts about 55 kg from breeding to farrowing.

## Housing management of pigs

Adequate housing and equipment for raising pigs are necessary to provide shelter against inclement weather, prevent diseases, control parasites and save labour.

The normal requirement of floor area, water and air space in pens for various classes of pigs is given below

Class of animals	Covered floor area per animal (m <sup>2</sup> )	Open-yard area per animal (m <sup>2</sup> )	Water required (litres)
Boar	6.25-7.5	8.8-12.0	45.5
Farrowing	7.5-9.0	8.8-12	18-22
Weaner	0.96-1.8	8.8-12	3.5-4
Dry sow	1.8-2.7	1.4-1.8	4.5-5

## **Creep space**

The flooring should have a rough finish and should be of a regular masonry type made up of water proof cement mortar. Proper drains should be provided so that the effluents are disposed off. Generally under village conditions the housing can be made up of pens measuring 3 m X 2.4 m or 3 m X 3 m with an open yard of nearly the same dimension or in some cases slightly longer. Walls should be 1.2-1.5 m high from the floor. For the purposes of farrowing some of the pens could be converted into farrowing pens by providing guard rails made up of G.I pipes of 5 cm diameter, along the walls, 20-25 cm from the ground and the wall. In addition to guard rails, creep space can be provided for the piglets along the wall by making a partition or in one of the corners with separate entrances for the piglets. This space usually of 0.75 m X 2.4 m area. In many of the farms the yard is provided with regular flooring.



Prolonged exposure of exotic breeds of pigs to bright sunshine may cause pigs to become overheated even during moderate weather. Shade helps in preventing deaths and increasing production efficiency during hot weather. While it is desirable to plant trees in the neighbourhood of pens for reducing the intensity of heat. But it is not desirable to plant trees for giving regular shade because they permit rapid build up of parasite levels.

## **Farrowing Pen**



## **Wallows**

Pigs have very few sweat glands. In areas having warm weather mature breeding animals and fattening animals need a wallow during summer months. Instead of permitting unsanitary wallows a masonry wallow with proper drainage would be desirable. The size of the wallow will depend upon the number and size of the animals.

## **Prevention and control of pig diseases**

- All pigs should be vaccinated against swine fever at the age of 2-4 weeks. Breeding pigs should be tested for brucellosis and leptospirosis. As a routine measure all young pigs at the time of weaning should be inoculated against swine fever.
- Animals purchased for the farm should be purchased from disease free herds. Newly purchased animals should be isolated from the other animals in the farm for a period of three to four weeks. No visitor allowed visiting the farm. Those stys or pig houses cleared of the animals are kept empty for three to four weeks for destruction of microorganisms causing the disease.